



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$\therefore dF = k\sqrt{2g(y-x)}dx$, where k = width of slit, and F = flow of water.

$$\text{Whence } F = k\sqrt{2g} \int_0^y \sqrt{y-x} dx = \frac{2k\sqrt{2g}}{3} y^{\frac{3}{2}}.$$

Call V the volume of water in the box at any instant.

Then $\frac{dV}{dt} = \frac{2k\sqrt{2g}}{3} y^{\frac{3}{2}}$. But $V = aby$, where a and b are the dimensions

of base of box. $\therefore \frac{abdy}{dt} = \frac{2k\sqrt{2g}}{3} y^{\frac{3}{2}}$.

$$\text{From which } t = \frac{3ab}{2k} \int_n^m y^{-\frac{1}{2}} dy = \frac{ab}{k\sqrt{2g}} \left[\frac{1}{\sqrt{n}} - \frac{1}{\sqrt{m}} \right], m \text{ and } n \text{ being the}$$

depths of water at beginning and end of time of discharge.

If $n=0$, or the box is emptied, $t=\infty$.

If $m=\infty$, $t = \frac{ab}{k\sqrt{2g}n}$; or the time to empty a box of infinite depth to a finite depth is finite.

37. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland.

A thin board, of which the elements are given, is balanced at the center but inclined at an angle. A sphere of known dimensions is put directly above the point of suspension and liberated. Find the motion of the system. That is, find (a) the time until the sphere leaves the board, (b) the ultimate angular velocity of the board.

Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Take the horizontal line through the point making the greatest angle with the plane in its initial position as the axis of x , and the axis of y vertically downward through the same point. Let R and T be the normal and tangential reactions of the plane and sphere at any time t from the commencement of motion, θ and ϕ the angles of rotation of the sphere and of the plane, m , k , a the mass, radius of gyration, and radius of the sphere, and r = the distance the sphere has moved on the plane, and x and y the coordinates of the center of the sphere.

Resolving horizontally and vertically, and taking moments about the center of the sphere,

$$m \frac{d^2 x}{dt^2} = R \sin \phi - T \cos \phi \dots \dots \dots (1), \quad m \frac{d^2 y}{dt^2} = mg - R \cos \phi - T \sin \phi \dots \dots \dots (2).$$

$$mk^2 \frac{d^2 \theta}{dt^2} = Ta \dots \dots \dots (3). \quad \text{We also have } \theta = \frac{r}{a} + \phi \dots \dots \dots (4),$$

$$x = r \cos \phi + a \sin \phi \dots \dots \dots (5), \text{ and } y = r \sin \phi - a \cos \phi \dots \dots \dots (6).$$

Eliminating T from (1) and (2), $\sin\phi\frac{d^2x}{dt^2} - \cos\phi\frac{d^2y}{dt^2} = \frac{R}{m} - g\cos\phi \dots (7).$

Eliminating T and R from (1), (2), and (3),

$$\cos\phi\frac{d^2x}{dt^2} + \sin\phi\frac{d^2y}{dt^2} = g\sin\phi - \frac{k^2}{a}\frac{d^2\theta}{dt^2} \dots\dots\dots(8).$$

$$\text{From (4), } \frac{d^2\theta}{dt^2} = \frac{1}{a}\frac{d^2r}{dt^2} + \frac{d^2\phi}{dt^2} \dots\dots\dots(9).$$

$$\begin{aligned} \text{From (5) and (6), } \frac{d^2x}{dt^2} &= \cos\phi\frac{d^2r}{dt^2} - 2\sin\phi\frac{dr}{dt}\frac{d\phi}{dt} - r\cos\phi\frac{d\phi^2}{dt^2} \\ &\quad - r\sin\phi\frac{d^2\phi}{dt^2} - a\sin\phi\frac{d\phi^2}{dt^2} + a\cos\phi\frac{d^2\phi}{dt^2} \dots\dots\dots(10), \end{aligned}$$

$$\begin{aligned} \frac{d^2y}{dt^2} &= \sin\phi\frac{d^2r}{dt^2} + 2\cos\phi\frac{dr}{dt}\frac{d\phi}{dt} - r\sin\phi\frac{d\phi^2}{dt^2} \\ &\quad + r\cos\phi\frac{d^2\phi}{dt^2} + a\cos\phi\frac{d\phi^2}{dt^2} + a\sin\phi\frac{d^2\phi}{dt^2} \dots\dots\dots(11). \end{aligned}$$

Eliminating $\frac{d^2x}{dt^2}$, $\frac{d^2y}{dt^2}$, $\frac{d^2\phi}{dt^2}$ from (7) and (8),

$$2\frac{dr}{dt}\frac{d\phi}{dt} + r\frac{d^2\phi}{dt^2} + a\frac{d\phi^2}{dt^2} = g\cos\phi - \frac{R}{m} \dots\dots\dots(12),$$

$$\frac{a^2 + k^2}{a^2}\frac{d^2r}{dt^2} - r\frac{d\phi^2}{dt^2} + \frac{a^2 + k^2}{a^2}\frac{d^2\phi}{dt^2} = g\sin\phi \dots\dots\dots(13).$$

(12) and (13) seem to indicate that one more condition at least should be given.

PROBLEMS.

46. Proposed by H. C. WHITAKER, A. M., Ph. D., Professor of Mathematics, Manual Training School, Philadelphia, Pennsylvania.

“There was an old woman tossed up in a basket

Ninety times as high as the moon.”

Mother Goose.

Neglecting the resistance of the air, how long did it take the old lady to go up?

47. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland.

What is the focus of the convex surface of a plano-convex lens, index μ , which will converge parallel monochromatic rays to a given focus, the rays entering the lens on the plane side?